

Comment on "Duality of x and ψ in Quantum Mechanics" by A.E. Faraggi and M. Matone

J.L. Lucio⁽¹⁾ and M. Ruiz-Altaba⁽²⁾

⁽¹⁾ *Instituto de Física, Universidad de Guanajuato, León, Gto., México*

⁽²⁾ *Instituto de Física, UNAM, A.P. 20-364, 01000 México, D.F.*

Recently, Faraggi and Matone introduced [1] an equation "dual" to the time-independent Schrödinger equation

$$-\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} = [E - V(x)] \psi, \quad (1)$$

namely

$$\frac{\hbar^2}{2m} \frac{\partial^2 x}{\partial \psi^2} = [E - V(x)] \psi \left(\frac{\partial x}{\partial \psi} \right)^3, \quad (2)$$

which they derived [1] in a rather roundabout way with the help of a "prepotential" $\mathcal{F}(\psi)$. They went on to speculate on a possible dual quantization of space based on this formula, employing also this formalism to study duality transformations in the pilot-wave version of quantum mechanics [2].

We would like to point out that, regardless of the conceptual interest and difficulties of interpreting space in terms of a wave-function instead of the opposite, equation (2) can be derived very easily from the trivial observation that

$$\frac{d\psi}{dx} = \left[\frac{dx}{d\psi} \right]^{-1}, \quad (3)$$

and thus

$$\frac{d^2x}{d\psi^2} = -\frac{d^2\psi}{dx^2} \left(\frac{dx}{d\psi} \right)^3. \quad (4)$$

Plugging (4) into (1) gives (2) directly. A more interesting truly dual version of (1) should involve a dual potential $\tilde{V}(\psi)$; we have no idea how to derive such an equation, nor what its interpretation would be.

Acknowledgements

Work partially supported by CONACyT under contract 3979PE-9608 and Catedra Patrimonial de Excelencia Nivel II.

Bibliography

- [1] A.E. Faraggi and M. Matone, *Phys. Rev. Lett.* **78** (1997) 163; hep-th/9606063.
- [2] A.E. Faraggi and M. Matone, hep-th/9705108..